arXiv:1610.07919v1 [math.GT] 25 Oct 2016

ON SUBCRITICALLY STEIN FILLABLE 5-MANIFOLDS

FAN DING, HANSJÖRG GEIGES, AND GUANGJIAN ZHANG

Abstract. We make some elementary observations concerning subcritically Stein fillable contact structures on 5-manifolds. Specifically, we determine the diffeomorphism type of such contact manifolds in the case the fundamental group is finite cyclic, and we show that on the 5-sphere the standard contact structure is the unique subcritically fillable one. More generally, it is shown that subcritically fillable contact structures on simply connected 5-manifolds are determined by their underlying almost contact structure. Along the way, we discuss the homotopy classification of almost contact structures.

1. Introduction

A closed contact manifold (M,ξ) is said to be *Stein fillable* if it arises as the boundary of a compact Stein manifold (W,J), and the hyperplane field ξ in the tangent bundle TM is given as the complex tangencies $TM\cap J(TM)$. It is well known that a Stein manifold of dimension $2n$ has a handle decomposition with handles of index at most equal to n. The filling is called *subcritical*, if there are no handles of index n.

In this note we are concerned with topological and contact geometric aspects of subcritically Stein fillable contact 5-manifolds. The first result we want to discuss gives a uniqueness statement for the diffeomorphism type of such contact manifolds when it has a finite cyclic fundamental group. This extends a corresponding result for simply connected contact manifolds due to Bowden–Crowley–Stipsicz [2]. The main issue is one of simple homotopy theory, which in our examples can be addressed with results of Hambleton–Kreck [6] on 2-complexes and, as in [2], the Mazur–Wall theory of thickenings.

In order to state the result, we need to introduce certain model manifolds. Let $m\geq 2$ be an integer. Write L_m for the 3-dimensional lens space $L(m,1)$ with an open 3-disc removed. This space L_m can be obtained from a solid torus $S^1\times D^2$ by attaching a 2-handle along an $(m,-1)$-torus not in $\partial(S^1\times D^2)$.

Oriented D^3-bundles over L_m are classified by the second Stiefel–Whitney class w_2 (this standard fact will be elucidated in the proof of Theorem 1.1). Since $H^2(L_m;\mathbb{Z}_2)$ is trivial for m odd, and isomorphic to $\mathbb{Z}_2$ for m even, the only D^3-bundles are the product $L_m\times D^3$ and, for $m=2n$ even, the non-trivial bundle $L_{2n}\tilde{\times}D^3$. After rounding of corners, we may think of the total spaces of these bundles as manifolds with boundary.

Similarly, over S^2 we have the trivial S^3-bundle $S^2\times S^3$, and the non-trivial one $S^2\tilde{\times}S^3$. In [2, Proposition 7.4] it was shown that if (M,ξ) is a closed, *simply connected* 5-dimensional contact manifold admitting a subcritical Stein filling, then

2010 *Mathematics Subject Classification.* 53D35; 32Q28, 57M20, 57R17.

M is diffeomorphic to $\#_r S^2\times S^3$ if M is spin, and $S^2\tilde{\times}S^3\#_{r-1}S^2\times S^3$ if M is not spin, where $r=\operatorname{rank} H_2(M;\mathbb{Z})$.

Our first result extends this to finite cyclic fundamental groups.

Theorem 1.1. *Suppose that (M,ξ) is a closed, connected contact 5-manifold admitting a subcritical Stein filling, with $\pi_1(M)\cong\mathbb{Z}_m$ for some integer $m\geq 2$. Set $r=\operatorname{rank} H_2(M;\mathbb{Z})$.*

(i) *If $m=2n+1$ is odd, then M is diffeomorphic to*

$$\partial(L_{2n+1}\times D^3)\#_r(S^2\times S^3) \text{ or } \partial(L_{2n+1}\times D^3)\#(S^2\tilde{\times}S^3)\#_{r-1}(S^2\times S^3),$$

depending on whether M is spin or not.

(ii) *If $m=2n$ is even, then M is diffeomorphic to*

$$\partial(L_{2n}\times D^3)\#_r(S^2\times S^3)$$

if M is spin, and to

$$\partial(L_{2n}\tilde{\times}D^3)\#_r(S^2\times S^3) \text{ or } \partial(L_{2n}\times D^3)\#(S^2\tilde{\times}S^3)\#_{r-1}(S^2\times S^3)$$

when M is not spin.

The diffeomorphism type of the subcritical Stein filling is determined by M.

On any of these manifolds, each homotopy class of almost contact structures contains a subcritically Stein fillable contact structure.

Our second result is concerned with the contactomorphism classification of simply connected subcritically Stein fillable contact 5-manifolds in terms of the first Chern class. Because of its special interest, we first formulate the statement for the 5-sphere. This theorem confirms an expectation from [4, Section 6]. Its proof is based on the fundamental work of Cieliebak–Eliashberg [3] on Stein manifolds.

Theorem 1.2. *Any subcritically Stein fillable contact structure on S^5 is isotopic to the standard contact structure.*

More generally, we shall prove the following. Here, for (M,ξ) a contact manifold, $c_1(\xi)\in H^2(M;\mathbb{Z})$ denotes the first Chern class of ξ; recall that a contact structure carries a complex bundle structure, unique up to homotopy [5, Proposition 2.4.8].

Theorem 1.3. *Let (M_i,ξ_i), $i=1,2$, be two simply connected subcritically Stein fillable contact 5-manifolds. If there is an isomorphism $\phi\colon H^2(M_1;\mathbb{Z})\to H^2(M_2;\mathbb{Z})$ such that $\phi(c_1(\xi_1))=c_1(\xi_2)$, then (M_1,ξ_1) and (M_2,ξ_2) are contactomorphic.*

In [4, Theorem 4.8] this result had been proved under the additional assumption that the fillings contain no 1-handles.

2. Homotopy classification of almost contact structures

In this section we discuss the homotopy classification of almost contact structures on 5-manifolds, correcting a negligence in [5]. Likewise, we describe the classification of almost complex structures on 6-manifolds, correcting a similar oversight in [16]. These classification results are relevant to the proof of Theorem 1.1 and Theorem 1.3.

Let M be a compact (not necessarily closed), oriented 5-manifold. A choice of Riemannian metric on M, or equivalently a reduction of the structure group of the tangent bundle to $\mathrm{SO}(5)$, allows us to describe the tangent bundle TM in terms of a classifying map $f\colon M\to \mathrm{BSO}(5)$. Define the inclusion $\mathrm{U}(2)\subset\mathrm{SO}(5)$ by the

embedding $\mathbb{C}^2 \equiv \mathbb{R}^4 \times \{0\} \subset \mathbb{R}^5$. Any subgroup $\mathrm{G} \subset \mathrm{O}(5)$ acts on the space $V(5)$ of orthonormal 5-frames in $\mathbb{R}^\infty$, and this defines the universal bundles $V(5) \to \mathrm{BG} := V(5)/\mathrm{G}$. The quotient BG is the classifying space for G-bundles, see [17, Section A.2]. The inclusion $\mathrm{U}(2) \subset \mathrm{SO}(5)$ defines a fibration $p\colon\ \mathrm{BU}(2) \to \mathrm{BSO}(5)$ with fibre $F_5 := \mathrm{SO}(5)/\mathrm{U}(2)$.

An almost contact structure on M is a reduction of the structure group of TM from SO(5) to U(2), which amounts to a lift $\tilde{f}$ of the classifying map f:



The lifting condition $p \circ \tilde{f} = f$ is equivalent to saying that the map

$$M \ni x \longmapsto \sigma(x) := \big(x, \tilde{f}(x)\big) \in M \times V(5)/\mathrm{U}(2)$$

is a section of the induced bundle $E := f^*V(5)/U(2) = f^*\mathrm{BU}(2)$ over M with fibre F_5. This is the obstruction-theoretic setting of [14, Part III]. Notice that $f^*V(5)$ is the frame bundle of M.

From now on we shall interpret almost contact structures on M as sections σ of this bundle $E \to M$. Homotopy of almost contact structures means homotopy of such sections.

Lemma 2.1. *The* U(2)*-bundle* $\tilde{f}^*V(5) \to M$ *corresponding to the almost contact structure defined by* $\tilde{f}$ *equals the pull-back of the* U(2)*-bundle* $f^*V(5) \to E$ *under the map* $\sigma\colon\ M \to E$*.*

Proof. Write the two relevant universal bundles as

$$\pi_{\mathrm{SO}}\colon\ V(5) \to \mathrm{BSO}(5) \quad \text{and} \quad \pi_{\mathrm{U}}\colon\ V(5) \to \mathrm{BU}(2).$$

Then

$$f^*V(5) = \big\{(x, v) \in M \times V(5)\colon\ f(x) = \pi_{\mathrm{SO}}(v)\big\},$$

and the bundle projection $\pi_E\colon\ f^*V(5) \to E$ is given by

$$\pi_E(x, v) = \big(x, \pi_{\mathrm{U}}(v)\big) \in f^*\mathrm{BU}(2) = E.$$

Under σ this pulls back to

$$\big\{(x, w) \in M \times f^*V(5)\colon\ \sigma(x) = \pi_E(w)\big\},$$

with the obvious projection map to M. This space can be rewritten as

$$\big\{(x, v) \in M \times V(5)\colon\ \tilde{f}(x) = \pi_{\mathrm{U}}(v)\big\},$$

which is the total space of the bundle $\tilde{f}^*V(5) \to M$. □

The fibre F_5 of the bundle $E \to M$ is diffeomorphic to $\mathbb{C}\mathrm{P}^3$, see [5, Lemma 8.1.2 and Proposition 8.1.3]. From the homotopy exact sequence of the generalised Hopf fibration $S^1 \hookrightarrow S^7 \to \mathbb{C}\mathrm{P}^3$ one then sees that the homotopy groups $\pi_i(F_5)$ are trivial for $i = 0, 1, 3, 4, 5$, and $\pi_2(F_5) \cong \mathbb{Z}$.

Since the fibre F_5 is simply connected, it is in particular 2-simple in the sense of [14, §16.5], i.e. the fundamental group operates trivially on $\pi_2(F_5)$. Moreover, the structure group SO(5) of the bundle $E \to M$ is connected. From [14, §30.4] it then

follows that the bundle of coefficients over M whose fibre over x is the homotopy group $\pi_2(E_x) \cong \mathbb{Z}$ of the fibre E_x of E is actually a trivial bundle. This implies that the obstruction to extending a section of E over the 2-skeleton of M to the 3-skeleton is a cohomology class in $H^3(M;\mathbb{Z})$. Given two sections of $E \to M$ that are homotopic over the 1-skeleton, the obstruction to homotopy over the 2-skeleton lives in $H^2(M;\mathbb{Z})$. Similarly, the obstruction cocycles are simply integral chains.

The obstruction class for the existence of a section over the 3-skeleton can be identified with the third integral Stiefel–Whitney class $W_3(M)$, see [5, p. 370]. By the vanishing of the other relevant homotopy groups of F_5, this class is the only obstruction to the existence of an almost contact structure. Likewise, the only obstruction to homotopy of two almost contact structures is the primary difference class in $H^2(M;\mathbb{Z})$.

We can now formulate the homotopy classification of almost contact structures. Regarding an almost contact structure as a U(2)-bundle, we can sensibly speak of its first Chern class c_1, which is a homotopy invariant. In the following statement and its proof we allow ourselves to identify an almost contact structure with the homotopy class it represents. The k-skeleton of M will be denoted by $M^{(k)}$.

Proposition 2.2. *Let M be a compact, oriented 5-manifold with $W_3(M) = 0$. There is a free and transitive action of $H^2(M;\mathbb{Z})$ on the set $\mathcal{A}(M)$ of almost contact structures on M. Write $u * \sigma \in \mathcal{A}(M)$ for the image of $\sigma \in \mathcal{A}(M)$ under the action of $u \in H^2(M;\mathbb{Z})$. Then $c_1(u * \sigma) = c_1(\sigma) + 2u$.*

Proof. Fix a reference element $\sigma_0 \in \mathcal{A}(M)$. For any $u \in H^2(M;\mathbb{Z})$, by the first extension theorem [14, §37.2] of obstruction theory, we can find a section σ'_u of E over $M^{(3)}$ such that the primary difference class $d(\sigma_0, \sigma'_u) \in H^2(M;\mathbb{Z})$ equals u. Since the higher relevant homotopy groups of F_5 vanish, σ'_u can be extended to a section σ_u over all of M.

Given any other $\tau_u \in \mathcal{A}(M)$ with primary difference $d(\sigma_0, \tau_u) = u$, the addition formula [14, §36.6] implies $d(\sigma_u, \tau_u) = 0$, and hence that σ_u and τ_u are homotopic. Thus, σ_u denotes a well-defined homotopy class. This allows us to define a free and transitive action of $H^2(M;\mathbb{Z})$ on $\mathcal{A}(M)$ by

$$u * \sigma_v := \sigma_{u+v}.$$

It remains to prove the formula for the first Chern class. From the homotopy exact sequence of the universal bundles we have $\pi_2(\mathrm{BU}(2)) \cong \pi_1(\mathrm{U}(2)) \cong \mathbb{Z}$ and $\pi_2(\mathrm{BSO}(5)) \cong \pi_1(\mathrm{SO}(5)) \cong \mathbb{Z}_2$. The homotopy exact sequence of the bundle

$$\mathbb{C}\mathrm{P}^3 = F_5 \hookrightarrow \mathrm{BU}(2) \longrightarrow \mathrm{BSO}(5)$$

then gives us

$$\begin{array}{ccccccc} \pi_2(F_5) & \longrightarrow & \pi_2(\mathrm{BU}(2)) & \longrightarrow & \pi_2(\mathrm{BSO}(5)) & \longrightarrow & \pi_1(F_5) \\ \mathbb{Z} & \longrightarrow & \mathbb{Z} & \longrightarrow & \mathbb{Z}_2 & \longrightarrow & 0. \end{array}$$

It follows that the first homomorphism in this sequence is multiplication by 2.

The inclusion map $\iota\colon F_5 \to \mathrm{BU}(2)$ is covered by a bundle map of U(2)-bundles:

$$\begin{array}{ccc} \mathrm{SO}(5) & \longrightarrow & V(5) \\ \downarrow & & \downarrow \\ F_5 & \xrightarrow{\iota} & \mathrm{BU}(2) \end{array}$$

This means that the bundle $\mathrm{SO}(5) \to F_5$ may be regarded as the induced bundle $\iota^*V(5) \to F_5$. By the observation on the homomorphism $\pi_2(F_5) \to \pi_2(\mathrm{BU}(2))$, the homomorphism

$$\iota^*\colon\ \mathbb{Z} \cong H^2(\mathrm{BU}(2);\mathbb{Z}) \longrightarrow H^2(F_5;\mathbb{Z}) \cong \mathbb{Z}$$

is likewise multiplication by 2. Since $H^2(\mathrm{BU}(2);\mathbb{Z})$ is generated by the first Chern class, it follows that $c_1(\iota^*V(5))$ is twice a generator of $H^2(F_5;\mathbb{Z}) \cong \mathbb{Z}$. Choose a generator of $\pi_2(F_5) = H_2(F_5;\mathbb{Z}) \cong \mathbb{Z}$ and the corresponding dual generator of $H^2(F_5;\mathbb{Z}) \cong \mathbb{Z}$ — in other words, fix an identification of these groups with $\mathbb{Z}$ — in such a way that $c_1(\iota^*V(5)) = -2$.

By construction, we have the formula $d(\sigma, u*\sigma) = u$ for the difference class. We therefore need to show that

$$(*) \qquad c_1(\tau) - c_1(\sigma) = 2d(\sigma,\tau) \ \text{ for any } \sigma,\tau \in \mathcal{A}(M).$$

It suffices to prove this formula over the 2-skeleton $M^{(2)}$. Indeed, the inclusion $M^{(2)} \to M$ induces an injective homomorphism $H^2(M;\mathbb{Z}) \to H^2(M^{(2)};\mathbb{Z})$.

The bundle $E|_{M^{(1)}}$ is trivial; moreover, the fibre F_5 is simply connected. Thus, we may assume that the sections σ and τ are constant (and identical) over the 1-skeleton $M^{(1)}$.

Recall from [14, §36] the definition of the primary difference class $d(\sigma,\tau)$, represented by a cochain with values in $\pi_2(F_5)$. Any oriented 2-cell $\Delta \subset M^{(2)} \subset M$ is described by a characteristic map $\varphi_\Delta\colon\ D^2 \to M$ sending $\mathrm{Int}(D^2)$ homeomorphically onto Δ, and ∂D^2 into $M^{(1)}$. The section σ of the bundle $E \to M$ defines a section σ_Δ of the pull-back bundle $\varphi_\Delta^*E \to D^2$ via

$$\sigma_\Delta(x) := \big(x, \sigma\circ\varphi_\Delta(x)\big),$$

likewise for τ:

$$\begin{array}{ccc} D^2 \times F_5 \cong \varphi_\Delta^*E & \xrightarrow{\overline{\varphi}_\Delta} & E \\ \sigma_\Delta,\tau_\Delta \uparrow\downarrow & & \downarrow\uparrow \sigma,\tau \\ D^2 & \xrightarrow[\varphi_\Delta]{} & M \end{array}$$

Notice that the pull-back bundle over D^2 is trivial, and in the trivialisation $\varphi_\Delta^*E \cong D^2 \times F_5$ the sections $\sigma_\Delta, \tau_\Delta$ may be regarded as maps $D^2 \to F_5$. However, there is no *a priori* relation between this trivialisation and that of $E|_{M^{(1)}}$, so $\sigma_\Delta, \tau_\Delta$ coincide over ∂D^2, but they will not, in general, be constant along ∂D^2.

Write $\pi_\pm\colon\ S^2_\pm \to D^2$ for the projection of the upper and lower hemisphere of the 2-sphere, respectively, onto the equatorial disc. Then the class $d(\sigma,\tau)$ is represented by the cocycle whose value on Δ is the element of $\pi_2(F_5)$ given by the map

$$d(\sigma,\tau)(\Delta) = \begin{cases} \sigma_\Delta\circ\pi_+ & \text{on } S^2_+, \\ \tau_\Delta\circ\pi_- & \text{on } S^2_-. \end{cases}$$

Here, by slight abuse of notation, we do not distinguish between cocycles and the cohomology classes they represent. The sign convention for the difference class is the standard one as in [14, §33.4].

Over E we have the $\mathrm{U}(2)$-bundle $f^*V(5) \to E$, which we shall now denote by η. Our aim is to compute the difference $c_1(\tau) - c_1(\sigma)$, which by definition equals

$c_1(\tau^*\eta) - c_1(\sigma^*\eta)$. These Chern classes live in the cohomology of M with coefficients in the coefficient bundle $\eta(\pi_1)$ in the notation of [14, §30.2]. Since the structure group $\mathrm{U}(2)$ has abelian fundamental group $\pi_1(\mathrm{U}(2)) \cong \mathbb{Z}$, hence is 1-simple, and is connected, again by [14, §30.4] this coefficient bundle is trivial and we are simply dealing with integral cohomology classes. (This is of course well known.)

With $\overline{\varphi}_\Delta$ defined by the diagram above, the pull-back bundle $\overline{\varphi}_\Delta^*\eta = D^2 \times \iota^* V(5)$ restricts to a trivial bundle over either $\sigma_\Delta(D^2)$ and $\tau_\Delta(D^2)$, and we have sections of these bundles over $\sigma_\Delta(\partial D^2) = \tau_\Delta(\partial D^2)$, since $\sigma \circ \varphi_\Delta|_{\partial D^2} = \tau \circ \varphi_\Delta|_{\partial D^2}$ is a constant section of $E|_{M^{(1)}} \cong M^{(1)} \times F_5$. These sections define elements of $\pi_1(\mathrm{U}(2)) \cong \mathbb{Z}$, and the classes $c_1(\sigma), c_1(\tau)$ are represented by the cochains whose value on Δ is precisely that respective element.

It follows that $c_1(\sigma) - c_1(\tau)$ is represented by a cochain whose value on Δ is given by the first Chern class of the $\mathrm{U}(2)$-bundle $\iota^* V(5)$ over the 2-sphere $d(\sigma,\tau)(\Delta) \in \pi_2(F_5)$. Since $c_1(\iota^* V(5)) = -2$, this implies $(*)$. □

Remark 2.3. The proposition implies that in the absence of 2-torsion in $H^2(M;\mathbb{Z})$, almost contact structures are determined up to homotopy by the first Chern class. This result was obtained earlier by Mark Hamilton [7].

By completely analogous arguments, one can also prove the following homotopy classification of almost complex structures on 6-manifolds. Again, the third integral Stiefel–Whitney class is the only obstruction to the existence of an almost complex structure.

Proposition 2.4. *Let W be a compact, oriented 6-manifold with $W_3(W) = 0$. There is a free and transitive action of $H^2(W;\mathbb{Z})$ on the set $\mathcal{A}(W)$ of almost complex structures on W. Write $u * \sigma \in \mathcal{A}(W)$ for the image of $\sigma \in \mathcal{A}(W)$ under the action of $u \in H^2(W;\mathbb{Z})$. Then $c_1(u * \sigma) = c_1(\sigma) + 2u$.* □

3. Topology of subcritically Stein fillable 5-manifolds

Proof of Theorem 1.1. Let (M,ξ) be a closed, connected 5-dimensional contact manifold with finite cyclic fundamental group, admitting a subcritical Stein filling (W,J). This Stein filling is made up of handles of index at most two, so W is simple homotopy equivalent to a finite 2-complex, see [10, p. 7]. Dually, W can be obtained from M by attaching handles of index at least four; in particular, the inclusion $M \to W$ induces an isomorphism on fundamental groups.

For $m \geq 2$ an integer, we write X_m for the m-fold dunce cap. This complex is obtained by attaching a 2-disc D^2 to the circle S^1 with an attaching map $\partial D^2 \to S^1$ of degree m. Equivalently, X_m can be obtained as a quotient space of D^2 by identifying each point $x \in S^1 = \partial D^2$ with its rotate through an angle $2\pi/m$.

According to [6, Theorem 2.1], given two finite 2-complexes K, K' of the same Euler characteristic $\chi(K) = \chi(K')$ and any isomorphism $\pi_1(K) \to \pi_1(K')$, where the fundamental group is a finite subgroup of $\mathrm{SO}(3)$, there is a simple homotopy equivalence $K \to K'$ inducing the given isomorphism on fundamental groups. This implies that if $\pi_1(M)$ is the cyclic group of order m, the Stein filling W is simple homotopy equivalent to the 2-complex $X_m \vee_r S^2$, where $r := \chi(W) - 1$. Thus, W is a thickening of this 2-complex in the sense of [10] or [15].

Because of $6 \geq 2\cdot 2+1$, 6-dimensional thickenings of a 2-complex are in the stable range, and by [10, Lemma 11.29] or [15, Proposition 5.1], oriented thickenings of

$X_m \vee_r S^2$ are classified up to diffeomorphism by $[X_m \vee_r S^2, \mathrm{BSO}]$. This set of homotopy classes is isomorphic to

$$[X_m \vee_r S^2, K(\mathbb{Z}_2, 2)] \cong H^2(X_m \vee_r S^2; \mathbb{Z}_2),$$

since the homotopy groups $\pi_k(\mathrm{BSO})$ coincide with those of the Eilenberg–MacLane space $K(\mathbb{Z}_2, 2)$ for $k \leq 2$. The isomorphism

$$[X_m \vee_r S^2, \mathrm{BSO}] \cong H^2(X_m \vee_r S^2; \mathbb{Z}_2)$$

is given by the second Stiefel–Whitney class, for this obstruction class detects the non-trivial oriented $\mathbb{R}^\infty$-bundle over S^2; see [5, Lemma 8.2.5], for instance.

Write $\natural$ for the boundary connected sum of manifolds with boundary, and $S^2 \tilde{\times} D^4$ for the non-trivial D^4-bundle over S^2. In the case that $m = 2n+1$ is odd, we have $H^2(X_{2n+1}; \mathbb{Z}_2) = 0$, so a thickening of $X_{2n+1} \vee_r S^2$ is determined by the tangent bundle over each of the r 2-spheres being trivial or not. There is a well-known diffeomorphism

$$S^2 \tilde{\times} D^4 \natural S^2 \tilde{\times} D^4 \cong S^2 \tilde{\times} D^4 \natural S^2 \times D^4,$$

see [4, Proposition 4.7]; this diffeomorphism can also be derived from the argument we shall use presently in the case that m is even. It follows that W diffeomorphic to

$$(L_{2n+1} \times D^3) \natural_r (S^2 \times D^4) \text{ or } (L_{2n+1} \times D^3) \natural (S^2 \tilde{\times} D^4) \natural_{r-1} (S^2 \times D^4),$$

depending on whether W is spin or not. Since the inclusion $M \to W$ induces an isomorphism on $H^2(\,.\,; \mathbb{Z}_2)$, this proves part (i) of the proposition.

If $m = 2n$ is even, we have $H^2(X_{2n}; \mathbb{Z}_2) = \mathbb{Z}_2$, so there is now also a choice of two thickenings over X_{2n}. The same argument as before shows that W is diffeomorphic to

$$(L_{2n} \times D^3) \natural_r (S^2 \times D^4)$$

if W is spin, or, in the non-spin case, to one of the three manifolds

$$\begin{aligned} W_{1,0} &:= (L_{2n} \tilde{\times} D^3) \#_r (S^2 \times D^4), \\ W_{0,1} &:= (L_{2n} \times D^3) \# (S^2 \tilde{\times} D^4) \#_{r-1} (S^2 \times D^4), \\ W_{1,1} &:= (L_{2n} \tilde{\times} D^3) \# (S^2 \tilde{\times} D^4) \#_{r-1} (S^2 \times D^4). \end{aligned}$$

The manifolds $W_{1,0}$ and $W_{0,1}$ are not diffeomorphic, since there is no isomorphism $H^2(W_{1,0}; \mathbb{Z}) \to H^2(W_{0,1}; \mathbb{Z})$ of $\mathbb{Z}_{2n} \oplus \mathbb{Z}^r$ whose mod 2 reduction sends $w_2(W_{1,0})$ to $w_2(W_{0,1})$; the same argument applies to the boundaries of these manifolds.

We claim, however, that $W_{1,1}$ is diffeomorphic to $W_{0,1}$; it suffices to prove this for $r = 1$. Both manifolds are obtained from a 6-ball by first attaching a 1-handle to produce $S^1 \times D^5$, and then a couple of 2-handles $h_1, h_2 \cong D^2 \times D^4$. In order to obtain $W_{0,1}$ we attach h_1 by an attaching map $\varphi_1 \colon \partial D^2 \times D^4 \to \partial(S^1 \times D^5)$ that sends $\partial D^2 \times \{0\}$ to a $(2n, -1)$-torus knot on

$$S^1 \times \partial D^2 \times \{0\} \subset S^1 \times \partial D^2 \times D^3 \subset \partial(S^1 \times D^5),$$

extended to an embedding of $\partial D^2 \times D^4$ with trivial framing. The handle h_2 is attached by a map φ_2 sending $\partial D^2 \times \{0\}$ to a homotopically trivial circle in

$$S^1 \times \partial D^2 \times \{0\} \setminus \varphi_1\big(\partial D^2 \times D^4\big),$$

extended to an embedding of $\partial D^2 \times D^4$ with twisted framing, corresponding to the non-trivial element of $\pi_1(\mathrm{SO}(4)) = \mathbb{Z}_2$.

By sliding h_1 over h_2 we get a diffeomorphic manifold where the framing of the first handle is now also twisted, in other words, the manifold $W_{1,1}$. This proves part (ii) of the proposition.

Finally, we come to the statement about the existence of subcritically fillable contact structures. Let M be one of the 5-manifolds in (i) or (ii), and W the corresponding 6-manifold discussed in the course of proving this classification, with boundary M. This manifold W admits an almost complex structure, since $H^3(W;\mathbb{Z}) = 0$. By [3, Theorem 1.5], any almost complex structures on W is homotopic to a subcritical Stein structure. The homomorphism $H^2(W;\mathbb{Z}) \to H^2(M;\mathbb{Z})$ induced by inclusion is an isomorphism, so by Propositions 2.2 and 2.4 every homotopy class of almost contact structures on M contains a subcritically Stein fillable contact structure. □

Remark 3.1. (1) Let M be one of the 5-manifolds in Theorem 1.1 and ξ a subcritically Stein fillable contact structure as just described. Then, by [1, Theorem 5.3], any symplectically aspherical filling is homotopy equivalent to the corresponding W, and even diffeomorphic to W if the Whitehead group of $\mathbb{Z}_m$ vanishes, which by [12, Corollary 6.5] happens exactly for $m \in \{2, 3, 4, 6\}$.

(2) From [6, Theorem B] one can derive the following stabilisation result. Let (M, ξ) be a closed, connected contact 5-manifold with finite fundamental group, admitting a subcritical Stein filling (W_0, J_0). Then any subcritical Stein filling of any contact structure on $M \# S^2 \times S^3$ is simple homotopy equivalent to $W_0 \natural S^2 \times D^4$. Homotopically, the additional summand amounts to a one-point union with S^2.

4. Uniqueness of subcritically Stein fillable 5-manifolds

Proof of Theorem 1.2. Suppose that ξ is a contact structure on S^5 that admits a subcritical Stein filling (W, J). This means that there is a contactomorphism $f\colon (S^5, \xi) \to (\partial W, \xi_J)$, where ξ_J denotes the contact structure induced by J.

By [1, Theorem 1.2] or an earlier result of Oancea–Viterbo [13], see the discussion in [1, Section 3.3], the manifold W is a simply connected homology ball, and hence diffeomorphic to the standard ball D^6 by Proposition A on page 108 of [11]. Choose an orientation-preserving diffeomorphism $G\colon D^6 \to W$.

Recall that any diffeomorphism of S^5 can be extended to a diffeomorphism of D^6; this corresponds with the fact that there are no exotic 6-spheres, see [8] and [9, Corollary VIII.(5.6)]. Let $\Phi\colon D^6 \to D^6$ be an orientation-preserving diffeomorphism extending the diffeomorphism $\left(G|_{S^5}\right)^{-1} \circ f\colon S^5 \to S^5$. Notice that the diffeomorphism $F := G \circ \Phi\colon D^6 \to W$ restricts to f on S^5.

It follows that the subcritical Stein structure F^*J on D^6 induces the contact structure ξ on S^5. Write J_{st} for the standard complex structure on D^6 inducing the standard contact structure ξ_{st} on S^5. The two complex structures F^*J and J_{st} on D^6 are homotopic as almost complex structures. Thus, by [3, Theorem 15.14], the respective induced contact structures ξ and ξ_{st} on S^5 are isotopic. □

Proof of Theorem 1.3. Our aim will be to show that any closed, simply connected contact 5-manifold (M, ξ) that admits a subcritical Stein filling (W, J) also admits a subcritical Stein filling without 1-handles. Then the theorem follows from the corresponding result [4, Theorem 4.8], which made precisely this additional assumption on the absence of 1-handles.

By the proof of [2, Proposition 7.4], the filling W is diffeomorphic to $\natural_r(S^2 \times D^4)$ or $(S^2 \tilde{\times} D^4)\natural_r(S^2 \times D^4)$ for some non-negative integer r (determined by M); this also follows from [1, Theorem 1.5]. As shown in the proof of [4, Proposition 4.5], for any class $c \in H^2(W;\mathbb{Z})$ that reduces modulo 2 to the Stiefel–Whitney class $w_2(W)$, there is a subcritical Stein structure on W *without* 1-*handles* with first Chern class c. (Moreover, the cited proposition shows that the contact structure induced on the boundary is determined by c.)

In particular, we find such a subcritical Stein structure J' with $c_1(J') = c_1(J)$. Since W is simply connected, our arguments in Section 2 show that J and J' are homotopic as almost complex structures. By [3, Theorem 15.14] this implies that J and J' are actually Stein homotopic. Thus, as claimed, the stipulation that there be no 1-handles poses no restriction. □

Acknowledgements. F. D. would like to thank Yakov Eliashberg and Otto van Koert for helpful conversations. F. D. and G. Z. are supported by grant no. 11371033 of the National Natural Science Foundation of China.

References

[1] K. Barth, H. Geiges and K. Zehmisch, The diffeomorphism type of symplectic fillings, `arXiv:1607.03310`.

[2] J. Bowden, D. Crowley and A. I. Stipsicz, The topology of Stein fillable manifolds in high dimensions I, *Proc. London Math. Soc. (3)* **109** (2014), 1363–1401.

[3] K. Cieliebak and Ya. Eliashberg, *From Stein to Weinstein and Back – Symplectic geometry of affine complex manifolds*, Amer. Math. Soc. Colloq. Publ. **59** (American Mathematical Society, Providence, RI, 2012).

[4] F. Ding, H. Geiges and O. van Koert, Diagrams for contact 5-manifolds, *J. London Math. Soc. (2)* **86** (2012), 657–682.

[5] H. Geiges, *An Introduction to Contact Topology*, Cambridge Stud. Adv. Math. **109** (Cambridge University Press, Cambridge, 2008).

[6] I. Hambleton and M. Kreck, Cancellation of lattices and finite two-complexes, *J. Reine Angew. Math.* **442** (1993), 91–109.

[7] M. Hamilton, On symplectic 4-manifolds and contact 5-manifolds, Ph.D. thesis, LMU München (2008); available at `https://edoc.ub.uni-muenchen.de/8779/`.

[8] M. A. Kervaire and J. W. Milnor, Groups of homotopy spheres I, *Ann. of Math. (2)* **77** (1963), 504–537.

[9] A. A. Kosinski, *Differential Manifolds*, Pure Appl. Math. **138** (Academic Press, Boston, MA, 1993).

[10] B. Mazur, Differential topology from the point of view of simple homotopy theory, *Inst. Hautes Études Sci. Publ. Math.* **15** (1963), 5–93.

[11] J. Milnor, *Lectures on the h-Cobordism Theorem* (Princeton University Press, 1965).

[12] J. Milnor, Whitehead torsion, *Bull. Amer. Math. Soc.* **72** (1966), 358–426.

[13] A. Oancea and C. Viterbo, On the topology of fillings of contact manifolds and applications, *Comment. Math. Helv.* **87** (2012), 41–69.

[14] N. Steenrod, *The Topology of Fibre Bundles*, Princeton Math. Ser. **14** (Princeton University Press, 1951).

[15] C. T. C. Wall, Classification problems in differential topology – IV. Thickenings, *Topology* **5** (1966), 73–94.

[16] C. T. C. Wall, Classification problems in differential topology – V. On certain 6-manifolds, *Invent. Math.* **1** (1966), 355–374.

[17] G. W. Whitehead, *Elements of Homotopy Theory*, Grad. Texts in Math. **61** (Springer-Verlag, Berlin, 1978).

School of Mathematical Sciences and LMAM, Peking University, Beijing 100871, P. R. China
E-mail address: dingfan@math.pku.edu.cn

Mathematisches Institut, Universität zu Köln, Weyertal 86–90, 50931 Köln, Germany
E-mail address: geiges@math.uni-koeln.de

School of Mathematical Sciences, Peking University, Beijing 100871, P. R. China
E-mail address: zhangguangjian8888@163.com